Uttar Pradesh: Horticultural Diversity





Brinjal or egg plant (*Solanum melongena* L.) is an important Solanaceous crop of sub-tropics and tropics. The name egg plant has been derived from the shape of the fruit of some varieties, which are white and resemble in shape to chicken eggs. The brinjal is of much importance in the warm areas of far East, being grown extensively in India, Bangladesh, Pakistan, China and the Philippines.

In India, it is one of the most common, popular and principal vegetable crops grown throughout the country except higher altitudes. It is a versatile crop adapted to different agro-climatic regions and can be grown throughout the year. It is a perennial but grown commercially as an annual crop. A number of cultivars are grown in India, consumer preference being dependent upon fruit color, size and shape. The varieties of *Solanum melongena* L. display a wide range of fruit shapes and colours. Most of the commercially important varieties have been selected from the long established types of the tropical India and China.



HYDERABAD INDIA 2012





Uttar Pradesh State Biodiversity Board

Ram Nagar Giant a local land race of Brinjal

Botanical Name: Solanum melongena L.; Candidate Varieties Name: Ram Nagar Giant; Varieties Type (Inbred/Hybrid): Inbred; Institution at which developed: Local Cultivar of Varanasi; Seasons (Kharif/Rabi/Zaid): Kharif.

Seedling: anthocyanin colouration of hypocotyl Stem: anthocyanin colouration Absent Stem: pubescence weak Leaf: length large (>20 cm) Leaf: width medium (10-20 cm) Leaf: blistering Present Leaf: spininess absent Leaf: louder of blade medium Leaf: colour of vein purple Leaf: intensity of colour of veins medium Inflorescence: number of flowers 1 to 3 Flower: size large Flower: colour White Flowering Time (days after seed sowing) Fruit: length medium (10-20 cm) Fruit: length / diameter ratio medium (1,0-2,0) Fruit: general shape ovoid Fruit: diameter of pistil scar small (<1,0cm)	S.No.Characteristics States				
3 Stem: pubescence weak 4 Leaf: length large (>20 cm) 5 Leaf: width medium (10-20 cm) 6 Leaf: margin entire 7 Leaf: blistering Present 8 Leaf: spininess absent 9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	1	9	absent		
4 Leaf: length (>20 cm) 5 Leaf: width medium (10-20 cm) 6 Leaf: margin entire 7 Leaf: blistering Present 8 Leaf: spininess absent 9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time (days after seed sowing) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	2	Stem: anthocyanin colouration	Absent		
(>20 cm) Leaf: width medium (10-20 cm) Leaf: blistering Present Leaf: spininess absent Leaf: lntensity of colour of blade medium Leaf: colour of vein purple Leaf: intensity of colour of veins medium Inflorescence: number of flowers 1 to 3 Helower: size large Flower: colour White Flowering Time medium (60-80 days) Fruit: length medium (10-20 cm) Fruit: length / diameter ratio medium (1.0-2.0) Fruit: general shape ovoid	3	Stem: pubescence	weak		
(10-20 cm) 6 Leaf: margin entire 7 Leaf: blistering Present 8 Leaf: spininess absent 9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time (days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	4	Leaf: length	· ·		
7 Leaf: blistering Present 8 Leaf: spininess absent 9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time (days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	5	Leaf: width			
8 Leaf: spininess absent 9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	6	Leaf: margin	entire		
9 Leaf: blade colour green 10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time (days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	7	Leaf: blistering	Present		
10 Leaf: Intensity of colour of blade medium 11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	8	Leaf: spininess	absent		
11 Leaf: colour of vein purple 12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	9	Leaf: blade colour	green		
12 Leaf: intensity of colour of veins medium 13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	10	Leaf: Intensity of colour of blade	medium		
13 Inflorescence: number of flowers 1 to 3 14 Flower: size large 15 Flower: colour White 16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	11	Leaf: colour of vein	purple		
14Flower: sizelarge15Flower: colourWhite16Flowering Time (days after seed sowing)medium (60-80 days)17Fruit: lengthmedium (10-20 cm)18Fruit: diameterLarge (>10 cm)19Fruit: length / diameter ratiomedium (1.0-2.0)20Fruit: general shapeovoid	12	Leaf: intensity of colour of veins	medium		
15 Flower: colour White 16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	13	Inflorescence: number of flowers	1 to 3		
16 Flowering Time medium (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	14	Flower: size	large		
(days after seed sowing) (60-80 days) 17 Fruit: length medium (10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	15	Flower: colour White			
(10-20 cm) 18 Fruit: diameter Large (>10 cm) 19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	16	<u> </u>			
19 Fruit: length / diameter ratio medium (1.0-2.0) 20 Fruit: general shape ovoid	17	Fruit: length			
(1.0-2.0) 20 Fruit: general shape ovoid	18	Fruit: diameter	Large (>10 cm)		
J 1	19	Fruit: length / diameter ratio			
21 Fruit: diameter of pistil scar small (<1.0cm)	20	Fruit: general shape	ovoid		
	21	Fruit: diameter of pistil scar	small (<1.0cm)		

S.No	.Characteristics	States
22	Fruit: shape of apex	indented
23	Fruit: colour of skin at commercial harvesting	greenish white
24	Fruit: intensity of purple colour of skin	light
25	Fruit: intensity of green colour of skin	light
26	Fruit: stripes	absent
27	Fruit: patches	absent
28	Fruit: glossiness at harvest maturity	weak
29	Fruit: size of calyx	medium
30	Fruit: colour of calyx	green
31	Fruit: intensity of colour of calyx	medium
32	Fruit: spininess of calyx	absent
33	Fruit: ribs	absent
34	Fruit: creasing of calyx	medium
35	Fruit: colour of flesh	greenish
36	Fruit: length of peduncle	medium (1.0-5.0 cm)
37	Fruiting: pattern	solitary
38	Plant: growth habit	erect
39	Plant: height	tall (101-150 cm)
40	Plant: spread (distance between two extremes leaf tips at widest point	medium (50-100 cm)
41	Fruit: color of skin at maturity	yellow

Special Features:

This cultivar is local selection from Varanasi. Plant habit is erect with dark green and broad foliage, 125-150 cm plant height with stout and 50% flowering at 60-70 days after sowing. It bears 3-4 fruits with light green colour and fruit weight ranged from 750g to 1500g. This cultivar is suitable for *bharta* making because of soft fleshed and few seeds.



Uttar Pradesh State Biodiversity Board

East Wing, III Floor, A-Block, PICUP Bhawan, Vibhuti Khand, Gomti Nagar, Lucknow-226 010, U.P., INDIA Phone: +91 522 4006746, +91 2306491, E-mail: upstatebiodiversityboard@gmail.com Website: www.upsbdb.org